

Avian Control™

is registered with the
US Environmental Protection Agency (EPA)
for use on these Crops*:

Apples

Blueberries

Cherries

Grapes

Strawberries

Raspberries

Blackberries

Sweet and Popping Corn

Sunflowers

Soybeans

Cereal Grains

Stone Fruit

Pome Fruit

(*see label for complete list and California Methods and Rates)

Carefully read and follow label directions.

Avian Control™

Drives birds away from barns,
packing sheds and other structures.

Avian Control™

Bird Problems Solved!



Avian Control™

**The Most Effective
Liquid Bird Repellent
Available.**

What experts and growers are saying about Avian Control™:

“Avian Control™ significantly reduced bird damage on my 1400 acres of sweet corn.”

— Taylor Vieck, JMR Farms

“Our experience using Avian Control™ on cherries in Washington State exceeded our expectations. The birds were gone!”

— Rob Valicoff, Valicoff Fruit Company, Inc.

“I had several Growers re-order in large quantities and said they have NEVER found anything that worked this well.”

— Ron Reister, Reister Grower Services

*“I made 3 bi-weekly applications at the low rate (32 ounces per acre) of Avian Control™ to several Rabbit Eye blueberry plants in June of 2012 in Raleigh, NC. Several nearby blueberry bushes did not receive any Avian Control™. Those that received no Avian Control™ had extensive bird damage with small pecks (over 40 percent) in the fruit and major fruit drop (over 75 percent). **Those blueberry bushes receiving Avian Control™ had very few pecks in the fruit (less than 5 percent) and with fruit drop of less than 5 percent.**”*

— Dr. Kenneth A Sorensen, Professor Emeritus
Department of Entomology
North Carolina State University

Recent and Ongoing Testing Conducted by:

- National Sunflower Association
- Penn State University
- Alicel Feed and Seed
- Outer Coastal Plain Vineyard Association
- North Carolina State University
- University of California-Irvine



Avian Control™

Features & Benefits

Features

- Effective On All Destructive Pest Birds
- Lasts Up To 14 Days
- Food Grade Formula - All Ingredients are “Generally Regarded as Safe” by the US Food and Drug Administration
- EPA Registered
- No PHI (Pre-Harvest Interval)
- Short 4 Hour REI (Re-Entry Interval)
- Ground or Aerial Spraying
- No Phytotoxicity

Benefits

- Stops Birds From Destroying Your Crops
- Affordable - Low Cost Per Acre
- Non-lethal, Suitable For Use Around People, Animals And Approved Crops.
- May Be Used Up To The Day Of Harvest
- Quick Re-entry Into Fields
- Will Not Alter The Taste of Treated Crops, Juice Or Wine
- Does Not Damage Plants or Foilage

Frequently Asked Questions

Q. What Makes Avian Control™ different from other bird repellents?

The active ingredient in Avian Control™ is Methyl Anthranilate. Avian Control™ does not harm birds, and unlike other products with Methyl Anthranilate, it has a strong molecular structure, making it a more effective and longer lasting repellent. It is also resistant to both biological and ultraviolet degradation and, in addition, it marks the treated crops with an infrared stain that alerts approaching birds that the crop has been treated.

Q. What types of birds does Avian Control™ work on?

The active ingredient in Avian Control™ has been demonstrated in numerous third-party field tests and university studies to repel nuisance birds such as starlings, blackbirds and grackles. All birds share a common pain nerve and Avian Control™ is designed to irritate this nerve when birds come into contact with the product through inhalation, preening, ingestion and even through their feet. Birds can never develop a tolerance to Avian Control™, insuring that the product will continue to be effective in the future.

Q. How much Avian Control™ do I use and how do I apply it?

Apply Avian Control™ at the rate of 12 ounces to 42 ounces per acre. Set your air blast or boom sprayer to apply 20 to 100 gallons of water per acre (a light mist) and make sure that your holding tank has internal agitation. Avian Control™ should be applied only to plants and crops. For aerial spraying consult the label for application rates.

Q. When should I start applying Avian Control™?

Apply Avian Control™ at the first sign of birds or bird damage.

Q. Can I mix Avian Control™ with other agricultural chemicals?

Avian Control™ may be mixed with other agricultural chemicals but, due to the possibility of chemical incompatibility, the effectiveness of Avian Control™ may be reduced. Avian Control™ is best applied alone.

Q. How long does Avian Control™ last and will rain wash it away?

Avian Control™ can last from 10 to 14 days depending upon the frequency and severity of local rain. Experience has shown that an application interval of 10 days is ideal for most crops. Avian Control™ is resistant to irrigation and light to moderate rain. Very heavy rains may wash Avian Control™ off of your plants and will require re-application.

Carefully read and follow label directions.