ROBERT MARVEL PLASTIC MULCH, LLC
2425 HORSESHOE PIKE
ANNVILLE, PA 17003
1-800-478-2214
717-838-0976 TEL
717-838-0978 FAX
info@robertmarvel.com
www.robertmarvel.com

Avian Control

Spraying Recommendations

Backpack or Electric Sprayers

1/8 acre (5,445 sq. ft.)	Mix 3 – 4 ounces of Avian Control with 1 – 1.5 ounces of liquid dish detergent (Dove, Palmolive or similar) and 1 – 2 gallons of water.
1/4 acre (10,890 sq. ft.)	Mix 6 – 8 ounces of Avian Control with 2 – 3 ounces of liquid dish detergent (Dove, Palmolive or similar) and 2 – 4 gallons of water.
1/2 acre (21,780 sq. ft.)	Mix 12 – 16 ounces of Avian Control with 4 – 6 ounces of liquid dish detergent (Dove, Palmolive or similar) and 4 – 8 gallons of water.
1 acre (43,560 sq. ft.)	Mix 24 – 32 ounces of Avian Control with 8 – 12 ounces of liquid dish detergent (Dove, Palmolive or similar) and 8 – 16 gallons of water.

Air Blast or Boom Sprayers

Apply according to label instructions using 24 – 32 ounces of Avian Control per acre with 25 – 100 gallons of water.

Tank Mixing

Avian Control may be mixed with other agricultural chemicals. Prior to mixing, blend a small amount of Avian Control with the other chemical in a jar of water to determine if they are compatible.

DO NOT mix Avian Control with chemicals or water with a pH below 7.0 (acidic) as this may reduce the effectiveness of Avian Control.

Frequency: Apply every 10 – 14 days. A heavy rain may require a new application sooner.

Inside Structures: Add 4 – 6 ounces of Avian Control to 2 gallons of water.

Initial Applications: Use the largest recommended amount of Avian Control.

Aggressive birds may require the concentration of Avian Control to be increased to 35 – 40 ounces per acre and the frequency to every 6 -8 days until they are “trained” to avoid sprayed areas.

Tip: Avian Control is most effective if applied before a severe infestation develops.